

Aflatoxins in Animal and Human Health

J.F. Robens* and J.L. Richard**

Contents

I. Introduction	69
A. Brief Historical Overview	70
II. Human Aflatoxicosis	71
A. Evidence from Analysis of Mutations	76
III. Aflatoxicosis of Animals	77
A. Swine	78
B. Cattle	78
C. Poultry	79
D. Carcinogenicity in Animals	80
E. Aflatoxin Interactions	81
F. Immunosuppression	82
Summary	84
References	85

I. Introduction

Although aflatoxins have been the cause of numerous deaths in livestock and suspected in some deaths of humans, the impact of aflatoxin consumption by a wide variety of animal species, including humans, extends much further than such a focal point. The economic significance of this group of toxins occurring in feeds and foods is through lowered productivity including meat and eggs, reduced weight gain, reduced feed efficiency, increased incidence of disease because of immunosuppression, and subtle damage to vital organs. Several human diseases have been recognized as being caused by aflatoxins, especially acute aflatoxicosis associated with highly contaminated foodstuffs (Shank 1977; Krishnamachari et al. 1975a, 1975b; Ngindu et al. 1982).

The modes of action and multiplicity of effects in humans and other animals are sufficiently complex, and in some cases obscure, to adequately provide prophylactic or therapeutic measures for aflatoxicosis. The intent of this overview is to provide the reader with an understanding of these effects

* USDA, ARS, NPS, APHS, Building 005, Barc-West, Room 210, Beltsville, MD 20705.

**National Center for Agricultural Utilization Research, 1815 North University Street, Peoria, IL 61604-3999.

based on experimental investigations or naturally occurring disease in humans and animals caused, at least in part, by aflatoxins.

A. Brief Historical Overview

The toxicity of a group of closely related compounds, called aflatoxin in the popular literature and often in the scientific press, was recognized in 1960 following one of the most intensive and productive investigations of unknown toxic factors of natural origin ever pursued. Peanut meal imported to Great Britain from Brazil was quickly targeted as the cause of the deaths of large numbers of turkey poults and ducklings on British farms. Mold hyphae were observed during examination for the presence of poisonous plants, although attempts at culture were negative (Goldblatt 1969). The disease syndrome was subsequently recognized in other domestic animals and in places other than England. The causative mold, *Aspergillus flavus*, that reproduced the animal disease signs and symptoms was actually isolated from meal associated with a hepatic problem occurring in ducklings in Uganda (Sargeant et al. 1961). The early history of the turkey outbreak in Great Britain was described by Blount (1960, 1961) and toxicity occurring in several farm species summarized by Allcroft (1965).

The identification of aflatoxins first utilized a bioassay with the duckling, a highly susceptible species, in which the proliferation of bile duct epithelial cells was clearly visible within a few days of the ingestion of toxic meal. It was soon discovered that the extraction of toxic meal yielded a highly fluorescent material that could be estimated visually and corresponded with the toxicity of the samples as determined biologically (Goldblatt 1969). The isolation and characterization of the four closely related toxins, aflatoxin B₁, B₂, G₁, and G₂, were first reported by Hartley et al. (1963), but their structures were not defined until 1965 by Buche and coworkers (Asao et al. 1965).

In early studies in 1961 with laboratory rats, liver tumors were noted (Lancaster et al. 1961) and came to the attention of the World Health Organization (FAO/UNICEF) who were considering peanut meal as a protein supplement in foods for undernourished children. Max Milner of UNICEF convened a meeting of interested scientists in New York City in October 1962 to help assess the hazards of aflatoxin to human health (Stoloff 1977; Wilson 1978). Also in the United States, an outbreak of trout hepatoma was recognized in the Spring of 1960 that dramatized and focused interest on the practical problems (Kraybill and Shimkin 1964), thus helping to set the stage for the extensive protracted search for the true role of aflatoxins in human liver disease (Goldblatt 1969).

W.H. Butler, in the proceedings of a symposium held at MIT in 1964, prophetically stated that "the carcinogenic properties of aflatoxin are now well documented, that large-scale epidemiological studies will probably be

required to demonstrate the role which these natural toxins play in the etiology of liver cancer," and, finally, "that the presence of traces of compounds such as mycotoxins in food warrants considerable attention as a health hazard to man." However, Oettle (1965) is credited with being the first investigator to draw serious attention to the fact that aflatoxin ingestion might cause human liver cancer.

II. Human Aflatoxicosis

Supporting the concern for the possible carcinogenicity of aflatoxin was the fact that the rat and rainbow trout were found to be highly susceptible to aflatoxin tumorigenesis. However, the excretion pattern of the rhesus monkey was found to be different from that of the rat, and results in primates were scarce and only proved they were much less susceptible to aflatoxin tumorigenesis than rats. Adamson et al. (1973) initiated a mixed sex and species study involving 40 rhesus and cynomolgus monkeys. By 44 months of age, one female who had received up to 800 µg/kg three times every 2 weeks became progressively jaundiced and inactive, was killed, and primary liver carcinoma was observed. Campbell and Stoloff (1974) collated the primate studies, and their discussion emphasized the variability in results of these studies. (See also the section on carcinogenicity in animals.)

In 1966–67, prospective human epidemiology studies were initiated in Africa when Alpert and coworkers (1969) surveyed the incidence of primary hepatoma in Uganda and determined aflatoxin levels in food samples stored for consumption between harvests (Alpert et al. 1969). The frequency of aflatoxin contamination of market food samples was positively associated with liver cancer incidence in localized population groups.

Coincident with the considerations of the possible role of aflatoxins in hepatocellular carcinoma was an effort to tie it to other distressing human illnesses of unknown etiology, i.e., kwashiorkor and Reye's syndrome. There is no generally agreed on definition of kwashiorkor, but edema associated with hypoalbuminemia is universally accepted as the essential minimum criterion for diagnoses (Hendrickse 1984). Kwashiorkor was first thought to be caused by protein deficiency in the presence of relative carbohydrate excess. When this hypothesis became untenable, it was noted that animals receiving aflatoxin experienced several effects consistently found in kwashiorkor, i.e., hypoalbuminemia, fatty liver and immunosuppression, and that "the world distribution and seasonal fluctuations of mycotoxins contamination of food are remarkably similar to the geographical distribution and seasonal presence of kwashiorkor which tends to peak in the wet season in many areas" (Hendrickse et al. 1983).

For example, aflatoxin AFB₁ was detected from autopsy liver specimens in all 36 children who had kwashiorkor (Hendrickse 1985). However, Hen-

drickse (1985) concluded only "that our findings in these studies establish an association between aflatoxins and kwashiorkor. The nature of this association is obscure and is being investigated." Frustration from the failure of medicine to prevent and reliably treat kwashiorkor made the association with dietary aflatoxin easy to take to the next step of a cause-and-effect relationship.

Reye's syndrome, an encephalopathy with fatty degeneration of the liver, was described by Reye et al. (1963). Following scattered reports of aflatoxin in tissues of victims of Reye's syndrome, Ryan et al. (1979) using formalin fixed tissue found an extremely strong association between children dying with Reye's syndrome and aflatoxin isolation from liver. However, Rogan et al. (1985), following an investigation by the Centers for Disease Control, concluded that "we do not believe that most or many cases of Reye's syndrome in the U.S. are caused by aflatoxin exposure; right now we think it prudent to view reports of aflatoxin in the tissues as indicators of exposure of uncertain significance for etiology." In another search for associations, the Office of Rural Health in Georgia (Caster et al. 1986) reported that the consumption of large amounts of corn, rice, peanuts, and milk (food considered to be potentially high in aflatoxins) was significantly related to the mental retardation of children in one county having high levels of aflatoxins in the food supply. No such relationship was found in another county having trivial levels of aflatoxins (<20 ppb) in its diet.

Acute toxic liver injury in humans has been ascribed to aflatoxin. The most extensive and widely publicized case was an outbreak in Western India in October 1974, where unseasonal rains resulted in extensive mold damage to corn crops with extremely high concentrations of aflatoxin found to be 6.3–15.6 ppm in corn from affected households. The illnesses were characterized by jaundice, rapidly developing ascites, portal hypertension, and a high mortality rate. Dietary calculations and the analysis of contaminated samples showed that affected people could have consumed between 2 and 6 mg of aflatoxin daily over a period of a month. Extensive bile duct proliferation, periportal fibrosis, and occasional multinucleated giant cell, typical of aflatoxin-induced disease in animals, were found in the histopathologic examination of liver specimens (Krishnamachari et al. 1975b).

There may be inhalation exposure to aflatoxin among industrial and agricultural workers. The presence of aflatoxin in dust from contaminated corn has been established (Burg and Shotwell 1984; Burg et al. 1981; Silas et al. 1987). Because of the sporadic distribution of aflatoxin in corn and other contaminated commodities, it was difficult to accurately quantitate the contamination. In industrial settings, two Czechoslovakian chemical engineers died working on a method for sterilizing peanut meal (Dvorackova et al. 1981) and two British biochemists developed adenocarcinomas of the colon following exposure to purified aflatoxins (Deger 1976). Van Nieuwenhuize

et al. (1973) identified a number of cancer cases among a small group of workers in a peanut processing plant in The Netherlands and associated them with the finding of aflatoxins at 300–400 mg/kg in the peanut meal. Hayes et al. (1984) conducted a follow-up study of workers in this plant. The test subjects extracted oil from peanuts and linseed oil, whereas the controls worked in a nearby corn processing plant and, because of the handling equipment in use, probably had much less dust exposure. The observed mortality due to cancer was reported to be higher than expected for the aflatoxin-exposed workers throughout the entire study period. However, the tumors were at a variety of sites with no primary tumors of the liver, but there was a much greater than expected number of tumors of the respiratory system in the aflatoxin-exposed group, with seven such deaths. The extent to which the development of tumors at individual sites was associated with aflatoxin exposure could not be determined.

There are several widely cited epidemiological studies attempting to link hepatocellular carcinoma with high intakes of aflatoxin B₁ from the diet. In the early group of studies, aflatoxin intakes were sampled in populations of different geographical areas and the results correlated with PLC incidence in those areas. These studies were by Shank et al. (1972) in Thailand, Peers and Linsell (1973) in Kenya, Van Rensburg et al. (1974) in Mozambique, Peers et al. (1976) in Swaziland, and Van Rensburg et al. (1985) in Mozambique/Transkei. In each of these locations except Mozambique, multiple areas were studied and aflatoxin exposure assessed by the measurement of dietary intake. Intake was not verified by measuring metabolites, DNA, or protein adducts in the urine of subjects.

Problems preventing strict adherence to good protocol design in these studies are discussed by Wagstaff (1985, 1990). These include short-term migration of working-age males in Africa, nonavailability of subjects, and difficulties in carrying out reliable analyses. In Kenya (Peers and Linsell 1973), the male population over 15 yr of age was 29% less than the female population of that group and the only liver cancer cases recorded were those people who voluntarily presented themselves to a hospital or clinic. The three study groups were not comparable, e.g., there were differences in altitude of the respective regions, proximity to the hospital, economic and agricultural conditions and crops grown, and diets. In Thailand, snacks and foods in short supply in homes, such as meats, were not sampled. Difficulties in case finding were so great in the Thailand study that incidence data in rural study areas were actually taken from nearby urban areas. Hepatitis B virus (HBV) status was not determined in any of these studies, although the possible role of HBV infection was discussed by Van Rensburg et al. (1985). Thus, with the exception of this latter study from Mozambique, the thesis was that aflatoxin might be the primary causative factor in areas of high primary liver cancer (PLC) incidence and, in fact, all the stud-

ies did demonstrate a positive correlation between aflatoxin intake and PLC incidence. However, in the United States, Stoloff (1983) found a small age-adjusted excess PLC mortality ratio for the Southeast and he questioned whether this small excess could be attributed to aflatoxin.

The early retrospective epidemiology studies depended on the accuracy of dietary surveys and analyses to establish aflatoxin consumption in the various populations surveyed. In addition, they assumed no infectious complicating factors in the etiology of PLC. In the years since the round of epidemiology studies were initiated in 1970, important advances have been made both in (1) identifying alternative etiologies and (2) tracking aflatoxin consumption. Epidemiological studies can now be carried out on a more scientific basis, thus assuring more confidence in their results.

The initial finding leading to the development of alternative etiologies for PLC was that of Sherlock et al. (1970), who noted the high prevalence of hepatitis B surface antigen (HBsAg) in a clinical series of PLC patients. The evidence for the association of PLC and HBV, aflatoxin and other possible risk factors has been summarized by Wagstaff (1985) and Bassendine (1987). Of importance in evaluating the role of HBV are such factors as the ecologic association of PLC and HBV in diverse regions throughout the world and case control studies comparing the prevalence of HBV serologic markers in PLC cases and controls. Any serious epidemiological study of PLC must now include testing for the HBsAg antigen to allow a concurrent evaluation of HBV with other risk factors in the test populations.

In the early studies, aflatoxin ingestion had to be estimated by taking appropriate samples of the diet and then assaying them by a reliable method before the aflatoxin content changed. Understandably, there was much room for error in this procedure. Confirmation of aflatoxin ingestion has now been made possible by the elucidation of its metabolism and detoxification pathways, identification of adducts, and development of rapid, sensitive, and accurate methods for adduct identification summarized by Groopman et al. (1988). An early finding was that of Campbell et al. (1970), who reported that aflatoxin M₁ was detectable in the urine of people who consumed more than 30 mg AFB₁/d. Later, after it was recognized that reactive electrophilic epoxides of many xenobiotics can covalently react with various nucleophilic centers in macromolecules, like DNA, RNA, and protein, such a pathway was also recognized for aflatoxin. The first major AFB₁-DNA adduct was identified by Essigmann et al. (1977) in an *in vitro* system; however, its presence was soon confirmed *in vivo*. The half-life of this adduct is approximately 12 hr, which severely limits its usefulness in confirming aflatoxin ingestion. Autrup et al. (1983) reported the presence of a putative AFB₁-Guanine product in urine collected from individuals living in areas with high liver cancer risk and suspected exposure to aflatoxin.

Aflatoxin B₁ binds to proteins as well as to DNA. The reaction with one or more lysine residues in serum albumin accounts for more than half of the total binding to this protein (Sabbioni et al. 1987). This adduct is a completely modified aflatoxin structure retaining only the coumarin and cyclopentene rings of the parent compound. Protein adducts are significant in biological monitoring because they represent the integrated dose of aflatoxin received over many previous weeks. The average half-life of albumin in humans is 20 d; thus, an accumulated dose of aflatoxin will be present after dietary exposure has ceased (Groopman et al. 1988).

Monoclonal antibodies are successfully used as the basis for tests to determine aflatoxin in foods and biological samples. These fast, relatively inexpensive, and easily performed tests allow the practical monitoring of human exposure to aflatoxin in field epidemiological studies (Groopman and Donahue 1988 and Garner et al. 1985). Monoclonal antibody columns, coupled with HPLC, can quantify aflatoxin-DNA adducts in human urine samples obtained from environmentally exposed humans (Groopman et al. 1988). These tests require strict quality control standards, e.g., the use of an internal standard is necessary for meaningful results; otherwise, variability in the assay can be high. The usefulness of this sensitive test still depends on adequate sampling of the diet of all test groups and the constancy of this diet over time, prospectively and retrospectively, as required by the particular protocol. A monoclonal antibody-based method to measure M₁ in urine was used by Sun et al. (1986) and Zhu et al. (1987) to document and quantify human exposure. Zhu found good correlation with diet only in male subjects.

The International Agency for Research in Cancer (IARC) has made several reports on the carcinogenicity of aflatoxin starting in 1976. This first report stated that "studies of liver cancer incidence in relation to aflatoxin intake provide circumstantial evidence of a casual relationship." However, by 1987, another IARC working group concluded that there was "sufficient evidence that aflatoxin is a probable human carcinogen." A later group of epidemiological studies—Sun and Chu (1984), Yeh et al. (1989), and Peers et al. (1987)—which reported an aflatoxin effect independent of HBsAg prevalence, were available to the latter working group and presumably influenced their conclusion (Campbell et al. 1990). However, the Sun and Chu report contained no aflatoxin intake data and the Yeh report compared only one "high" and one "low" village in the Guangxi region of China. In the third study, Peers and coworkers reinvestigated four regions in Swaziland studied earlier and were the first to utilize both food measurements of aflatoxin intake and HBV exposure of the same population. However, they had to estimate the consumption of peanuts, which were by far the most contaminated foodstuff, and the incidence of primary liver cancer was surprisingly low, being surpassed by cervical and esophageal cancer.

In another recent study, Autrup et al. (1987) reported no aflatoxin effect on PLC incidence rates when all ethnic, social, and cultural groups were combined but a positive association was noted among the Bantu people. Autrup et al. (1987) also measured HBsAg carrier rates. Yeh et al. (1989) reported on their study of HBV status, aflatoxin exposure, and PLC incidence conducted in southern Guangxi China between July 1982 and June 1987. In a cohort of 7,917 men aged 25–64 yr, a total of 149 deaths occurred, 76 of which (51%) were due to PLC. Prevalence of HBsAg positivity in PLC deaths was 91%, in contrast to 23% of all members of the cohort. However, the authors saw no HBsAg effect in the cross-sectional case analysis. There was an almost linear relationship between estimated aflatoxin B₁ exposure in the subpopulations and corresponding mortality rates of PLC. However, only the average regional aflatoxin exposure values were available and not the individual exposure values.

The latest epidemiology study to be reported that examined the association of PLC with dietary aflatoxin, among other factors, is fortunately far and away the most comprehensive, with 6,500 participants. This study by Campbell et al. (1990) was a comprehensive cross-sectional survey in the People's Republic of China on the possible risk factors for primary liver cancer. It included 48 survey sites, an aflatoxin exposure range of about 600-fold, a 39-fold range of PLC mortality rates, and a 28-fold range of HBsAg+ carrier prevalence. Although the accuracy of the urinary assay used for measuring aflatoxin metabolites has been questioned, this study did estimate exposures for a large number of nutritional, dietary, and lifestyle variables. It represents a much larger number of survey sites, a wider range of PLC mortality rates, and a wider range of aflatoxin exposure than in the previous surveys. In addition, aflatoxin consumption was estimated by measuring aflatoxin metabolites in urine.

In Campbell's study, in contrast to the reported results of some of the previous studies, there was no association between aflatoxin exposure and PLC mortality. There was also a lack of association of PLC mortality with corn and moldy peanuts, the two foods in the diet commonly contaminated with aflatoxin. However, PLC mortality was positively correlated to the prevalence of HBsAg and other factors. Notably, in the United States, Stolf (1983) found a small age-adjusted excess PLC mortality ratio for the Southeast. However, there was serious question whether the excess PLC mortality could be attributed to aflatoxin.

A. Evidence from Analysis of Mutations

The interaction of chemicals or their metabolites with DNA is thought to be a major factor in chemical carcinogenesis. Tissue metabolism is important in the formation of nonreactive and reactive metabolites following exposure

to chemical toxicants. Some of the reactive metabolites formed in tissues can irreversibly bind to DNA. If the damaged DNA is not repaired and the cell subsequently divides, the damage can be fixed, resulting in a mutational event. Following mutation, a series of events is initiated, ultimately leading to an adverse biological effect such as cancer. This series of events is under investigation, with many suspect chemicals including aflatoxin.

In a recent series of papers in *Nature*, Hsu et al. (1991), Bressac et al. (1991), and Harris (1991) reported on the incidence of mutations in p53, a putative tumor suppressor gene in PLC from patients in the high-risk areas of southern Africa and China. The mutations in both areas were found to be clustered on codon 249 with transversions of G to C or to T. These base changes are produced by a range of carcinogens, including aflatoxin. Because the papers reported only on the work delineating the effect of aflatoxin on p53, they do not conclusively link the incidence of PLC with either suspected etiological agents.

The papers provide supporting evidence for the possibility that p53 mutations caused by aflatoxins or other environmental carcinogens might contribute to the high incidence of PLC in other areas. In the discussion of such an association with aflatoxin, the authors point out that the integration of HBV DNA is a frequent event associated with PLC. A few of these integrations induce insertional mutations in host genes, but are unlikely to cause point mutations, as observed in the reported work with aflatoxin. Before any definitive conclusions can be drawn, further work will be necessary, particularly on the effects of HBV in various hepatoma cell lines.

III. Aflatoxicosis of Animals

The major target organ for aflatoxins in animals is the liver. These compounds have been found to be carcinogenic and teratogenic in animals, as well as the cause of impairment of protein formation, coagulation, weight gain, and immunogenesis.

Domesticated animals are variably susceptible to aflatoxins; some oral LD₅₀ values are given for aflatoxin B₁ in several animal species in Table 1. Sheep and cattle appear to be more resistant than most livestock and poultry, whereas swine, chickens, turkeys, and ducklings are more susceptible (Pier 1981). Within a species, there are breed differences; this is best demonstrated in breeds of chickens (Dalvi 1986). Although these factors influence the toxic effects of aflatoxin, other factors to consider are age, sex, exposure (duration and dose), nutrition, environmental stresses, and other chemical exposures including other mycotoxins.

Table 1. Oral LD₅₀'s of Aflatoxin B₁

Animal	Sex	Age/Size	LD ₅₀ (mg/kg)
Duckling	M	1 d	0.37
Rat	M-F	1 d	1.0
Rat	M	21 d	5.5
Rat	F	21 d	7.4
Rat	F	150 g	17.9
Dog	M-F	Adult	ca. 0.5
Hamster	M	30 d	10.2

A. Swine

Liver changes in swine with acute aflatoxicosis include centrolobular congestion and hemorrhage with the hepatocyte nuclei showing the margination of chromatin and some pyknosis (Cysewski et al. 1968). These changes were followed by centrolobular necrosis and were concomitant with increases in serum enzymes (OCT and AST) and prothrombin time and decreased liver function as measured by bromosulphaphthalein clearance. Pigs so affected became lethargic, did not eat, and eventually became hypothermic, icteric, and had blood in their feces. Additional studies have demonstrated that the major change observed in swine aflatoxicosis is damage to the hepatic parenchyma (Panangala et al. 1986; Miller et al. 1981, 1982; Cook et al. 1989). Panangala and coworkers (1986) found that weanling swine fed 500 ppb dietary aflatoxin had reduced growth rate, feed efficiency, and general unthriftiness. They felt that feeds containing 300 ppb aflatoxins could affect growth rate with prolonged feeding. These studies agree with those of Duthie et al. (1966), Southern and Clawson (1979), and Sisk and Carlton (1972) and influenced the decision of the FDA in establishing 200 ppb as the guideline for aflatoxin in corn for finishing swine. The aflatoxins are not known to significantly affect reproduction in swine (Armbrecht et al. 1972; Hintz et al. 1967). However, as discussed later in this chapter, aflatoxins in sow's milk can affect piglets and *in utero* exposure can affect the integrity of the immune system in neonatal pigs.

B. Cattle

The acute intoxication of cattle with aflatoxins produces liver lesions similar to those described for swine, although cattle are less susceptible to the aflatoxins (Pier et al. 1976; Bodine and Mertens 1983). Affected animals may exhibit reduced feed consumption, weight loss, and reduction in milk production. Masri and coworkers (1969) found that dietary aflatoxin could induce a 33% decrease in milk yields of dairy cattle.

The more significant effects in cattle may occur when animals consume lower levels of aflatoxins and chronic effects appear such as reduced reproductivity, immunosuppression, and reduced feed efficiency. These effects are economically important to the producer, as demonstrated by Guthrie (1979) in which a dairy herd fed 120 ppb aflatoxin-contaminated corn had a 2% reduction in breeding efficiency over a 5-mon period and milk production increased 28% when the cows were given a noncontaminated diet. General health problems occurred in this herd, including small calves, diarrhea, mastitis, reduced feed consumption, and respiratory illness.

Dairy cattle convert dietary aflatoxin rather efficiently to aflatoxin M₁ that occurs in milk. Aflatoxin at 30 ppb in feed will result in milk residue levels of less than 1 ppb (Park and Pohland 1986; Price et al. 1984; Applebaum et al. 1982). Aflatoxin M₁ can occur at 1-hr postdosing in a dairy cow given 0.5 mg aflatoxin B₁/kg of body weight (Trucksess et al. 1983). Mertens (1979) determined that about 0.9% of ingested aflatoxin is converted to aflatoxin M₁ in milk. Similar results were obtained by Van Dijk and colleagues (1984). This metabolic product is slightly less toxic than aflatoxin B₁ (CAST 1989). However, Van Dijk and colleagues (1984) did not find physiologic or biochemical differences in neonatal dairy calves fed milk containing levels of aflatoxin M₁ approximating that which would occur naturally.

Because aflatoxin remained in the rumen content of steers after a 2.5 wk withdrawal period following their being fed approximately 460 ppb dietary aflatoxin for 15 wk, Richard and colleagues (1983) suspected that the rumen function may have decreased. When this hypothesis was subsequently tested using radiotelemetric procedures, they found that rumen motility was reduced in steers given single doses of aflatoxin (Cook et al. 1986). Earlier work had demonstrated that aflatoxins affected *in vitro* and *in vivo* rumen functions such as cellulose digestion, volatile fatty acid formation, and proteolysis (Fehr and Delage 1970; Dvorak et al. 1977).

Thus, the general effect of aflatoxin in cattle is liver disease, with specific effects ranging from acute to chronic disease dependent on the dosage given. Milk production in cows appears to be affected even by low levels of dietary aflatoxins that have no other effects on the health of the animals (Patterson and Anderson 1982). Aflatoxin M₁ can occur readily in the milk of dairy cows given dietary aflatoxin and can cause disease in calves consuming contaminated milk, providing there is sufficient level of contamination (Van Dijk et al. 1984). The FDA has established a 100-ppb guideline level for aflatoxin in corn for breeding cattle, but a 20-ppb limit for corn fed to dairy cattle (CAST 1989).

C. Poultry

The poultry industry is probably more severely affected than any of the livestock industries because poultry appear to be more susceptible to the effects of dietary aflatoxins and losses can be quite severe (Hamilton 1971).

Susceptibility to aflatoxins varies in chickens, depending on dosage, nutritional status, environmental stress, breed, strain, and sex. Low concentrations of dietary aflatoxins can cause weak chickens with reduced weight gains, feed efficiency, and egg production. The latter was noted in at least two studies (Garlich et al. 1973; Wolzak et al. 1985). Again, the primary damage is to the liver, which may be pale, mottled, and have petechial hemorrhages. There may be swelling and vacuolation of hepatocytes and bile duct proliferation. Aflatoxin is a probable cause of fatty liver syndrome in laying hens, a disease causing considerable loss in some large laying flocks (Hamilton and Garlich 1971). Concomitant with pathologic changes in the liver, there are increases in liver-specific serum enzymes.

Although turkeys appear to be more susceptible to aflatoxins than chickens, the effects of aflatoxins on this species are similar to those found in chickens (Richard et al. 1986). Mortalities can occur in poults due to acute hepatic necrosis and hemorrhage after consumption of 1 ppm dietary aflatoxin (Pier 1981).

Of all avian species tested, ducklings appear to be the most susceptible to aflatoxins and, for this reason, they are used in the bioassay for aflatoxins. Acute aflatoxicosis of ducklings is similar to that in turkeys and chickens.

Kratzer et al. (1968) reported that levels of dietary aflatoxin of 1,600 ppb fed for 8 wk did not produce detectable residues in the edible meat of broilers. Similarly, no detectable aflatoxin was found in the meat or eggs of laying hens fed 2,700 ppb aflatoxin for 48 d. In the same study, Rhode Island red chicks were found to be more susceptible than Arbor-Acres hybrid chicks.

D. Carcinogenicity in Animals

The toxicity and carcinogenicity of aflatoxin B₁ are influenced by a variety of factors including, among others, diet, microsomal activity, age, sex, and animal species. The recipient of the toxic insult metabolizes or activates the aflatoxins primarily by the microsomal mixed-function oxidase system of the liver (Shimada and Guengerich 1989). The parent compound is normally detoxified by conversion to hydroxylated intermediates (e.g., aflatoxins Q₁, M₁, and B_{2a}) that are water-soluble and eliminated via urine or bile. However, certain aflatoxin B₁ intermediates may be formed, such as the 8,9 epoxide (Baertschi et al. 1988), that react with macromolecules (DNA, RNA, and protein) and form adducts (Busby and Wogan 1981). The major DNA adduct formed both *in vivo* and *in vitro* in this reaction is 8,9-dihydro-8-(N7-guanyl)-9-hydroxy aflatoxin B₁ (Gopalakrishnan et al. 1990). These adducts are suspect in the toxicity, especially mutagenicity and carcinogenicity, of aflatoxins in animals (Busby and Wogan 1981). The carcinogenicity of aflatoxins has been demonstrated in a variety of animal species; ducks

given 0.035 ppm aflatoxin developed hepatic tumors in 14 mon (Carnaghan 1965). Trout hepatomas were demonstrated in 96% of the fish fed 20 ppb aflatoxin for 20 mon (Halver 1969). All male rats given 15 ppb dietary aflatoxin B₁ for 68 wk developed hepatocellular carcinomas, whereas the same was true for female rats fed the same diet for 80 wk (Wogan and Newberne 1967). Since these early studies, neoplasms have been induced by aflatoxins in other species such as ferrets, guppies, mice, monkeys, and sheep (CAST 1989). Shalkop and Armbrecht (1974) did report aflatoxin induced hepatic cell adenomas in brood sows fed aflatoxins for 20 mon. However, the importance of carcinogenicity in livestock is diminished because of the relatively short time that these animals are likely to be fed a contaminated diet prior to the time they are marketed for slaughter.

E. Aflatoxin Interactions

The probability that more than one toxin may interact in producing toxicity in animals is high for the following reasons:

1. Several toxigenic fungi often occur in the same batch of feed.
2. Some fungi are capable of simultaneously producing more than one mycotoxin in a substrate.
3. Multiple mycotoxins produced by different fungi have been found in feeds.

Most of the efforts to systematically study some of these interactions have been performed in poultry and certain laboratory animals, as well as swine. Combinations of aflatoxins and several other mycotoxins have been studied. Aflatoxin and T-2 toxin interacted synergistically when given in combination to broiler chicks at dietary concentrations of 2.5 and 4 µg/g, respectively (Huff et al. 1988), as demonstrated by decreases in body weight; increases in the relative weights of the kidney, gizzard, and heart; and decreases in mean corpuscular volume and serum potassium levels.

An important event occurred, with significant application to the diagnosis of aflatoxicosis, when aflatoxin and ochratoxin A were fed to broiler chickens (Huff and Doerr 1981). Although dietary aflatoxin at 2.5 µg/g caused lipid accumulation in the liver, a diagnostic feature of aflatoxicosis in poultry, this phenomenon was inhibited by the simultaneous feeding of 2.0 µg/g of ochratoxin A. This is significant when the diagnosis of aflatoxicosis is made based on signs of the disease (i.e., fatty liver) in poultry. Also, these concentrations of dietary mycotoxins were synergistic in producing increased kidney and gizzard weights relative to body weights. When the same concentrations of these toxins were tested in broilers for the effects on immunity, only complement activity was affected but was likely attributed to the aflatoxin only (Campbell et al. 1983). Similarly, the body weight depression of broilers by this toxin combination persisted longer after a with-

drawal period than did that induced by the individual toxins (Huff et al. 1983). Increased susceptibility to bruising was noted by these workers in birds given these toxins individually and in combination. These authors have studied aflatoxin in combination with deoxynivalenol (Huff et al. 1986) and found no synergistic activity in broilers in several measured parameters.

F. Immunosuppression

One of the major economic considerations of the aflatoxin problem is that of impairment of immunogenesis and native resistance in animals. Within the first decade of research following the recognition of aflatoxins, work had been published on the effect of these toxic metabolites on various immune phenomena (Richard et al. 1978).

One of the initial studies demonstrated that antibody formation in vaccinated mice was affected, whereby typhoid agglutinin titers were depressed (Galikeev et al. 1968). Neither agglutinins nor preceptins to *Pasteurella multocida* in turkeys fed aflatoxin B₁ during or after vaccination were affected (Pier and Heddleston 1970). Similar results were found in the hemagglutinins to *Salmonella pullorum*, as measured by an indirect hemagglutinin test, in chicks given aflatoxin (Adinarayanaiah et al. 1973). Aflatoxin did not affect the hemagglutination inhibition or serum neutralization titers to Newcastle disease virus in turkey poult (Pier et al. 1971). Guinea pig titers to *Brucella abortus* agglutinin titers were not affected by aflatoxin ingestion (Thurston et al. 1974). The depression of agglutinin titers to sheep erythrocytes was found in chicks dosed with aflatoxin (Thaxton et al. 1974). The type of antigen and involvement of cells may be the cause for variation in the antibody effects due to aflatoxin. More recently, studies demonstrated that antibody production was decreased when directed against T-dependent antigen and sheep red blood cells, but was unaffected in animals challenged by T-independent antigens, such as lipopolysaccharide (Reddy et al. 1987).

Certain nonspecific humoral factors associated with a variety of immune phenomena have been affected by aflatoxins and may be related to the ability of aflatoxins to interfere with protein synthesis. Complement activity was depressed in guinea pigs dosed with aflatoxins (Thurston et al. 1972). The activity of this serum component has been depressed by aflatoxin also in poultry, swine, and cattle (Richard 1991). Guinea pigs and cattle given aflatoxins also have a decreased bacteriostatic serum factor to *Escherichia coli* (Thurston et al. 1986). An unknown nonspecific humoral factor appeared to be affected in the aflatoxin-impaired acquired immunity to *P. multocida* in turkey poult (Pier et al. 1971).

The effects of aflatoxins on the resistance of animals to infectious disease agents are somewhat variable-dependent on organism, toxin dose and con-

Table 2. Aflatoxin-Induced Changes in Response of Animals to Infectious Disease Agents

Organism	Animal Species	Response	Reference
<i>Candida albicans</i>	Chickens	Increased susceptibility	Hamilton and Harris (1971)
<i>Aspergillus fumigatus</i>	Turkeys	No change	Richard et al. (1973)
<i>Eimeria tenella</i>	Chickens	Increased susceptibility	Edds et al. (1973) Wyatt et al. (1975)
<i>Salmonella gallinarum</i>	Chickens	No change	Smith et al. (1969)
<i>Salmonella</i> spp (4)	Chickens	Increased susceptibility	Boonchuvit et al (1975)
<i>Mycobacterium paratuberculosis</i>	Hamsters	No change	Larsen et al. (1975)
<i>Erysipelothrix rhusiopathiae</i>	Swine	Immunosuppression	Cysewski et al. (1978)
<i>Treponema hyodysenteriae</i>	Swine	Increased susceptibility	Joens et al. (1981)
<i>Streptococcus</i> sp and <i>Staphylococcus</i>	Bovine	Increased mammary inflammation	Brown et al. (1981)
Newcastle disease virus	Chicken	Increased susceptibility	Byong and Rothenbacker (1982)
Infectious bronchitis virus	Chicken	Increased susceptibility	Ratanesthankul (1983)

stitution, animal species, and perhaps sensitivity of the test. Table 2 lists the various parameters that have been tested in this regard. Doerr and colleagues (1987) found that chickens given aflatoxins were more sensitive to the virulence factors beta hemolysins of the pathogen *Staphylococcus*.

The likely importance of aflatoxins regarding immunity is that they affect the cell-mediated responses. These effects are apparently related to functional aspects of the involved cells, such as lymphokine production and the processing of antigens by macrophages (Pier and McLoughlin 1985). Aflatoxin-induced reduction in the capacity of macrophages to phagocytize particles has been demonstrated (Michael et al. 1973; Richard and Thurston 1975). Richard and Thurston (1975) found that a heat-stable serum substance important in phagocytosis was reduced in rabbits given aflatoxin orally for 3 wk.

The functional aspects of lymphocyte responses to mitogens and production of lymphokines involved in migration inhibition are reduced by aflatoxins (Pier 1986). Similarly, the graft vs host response was reduced in chickens given aflatoxin (Giambrone et al. 1978). Changes in T- and B-cell populations were not found in the peripheral blood of guinea pigs given aflatoxin (McLoughlin et al. 1984). Lymphocytes with HLA-A3 antigen had a lower natural killer activity toward target tumor cells and were more highly suppressed with aflatoxin B₁ than control cells when stimulated by phytohemagglutinin (Cheng-ya et al. 1987).

The proximal entity in a number of the effects of certain mycotoxins on animal systems is unknown, but differences in effects on the immune process among aflatoxin B₁ and its metabolites do exist (Bodine et al. 1983). Aflatoxin Q₁ was active in suppressing *in vitro* lymphoblastogenesis using bovine lymphocytes. Others have found similar results with porcine lymphocytes (Yang 1983).

Immunosuppression has been demonstrated in animals following embryonic exposure. Aflatoxin was transferred across the porcine placenta, and the *in utero* exposed piglets sensitized with killed *Mycobacterium paratuberculosis* had a reduced delayed cutaneous hypersensitivity when skin-tested with johnin 2 wk later (Pier et al. 1985). When chicken embryos were dosed with 0.1 µg of aflatoxin B₁, the graft vs host response was reduced when assayed with a splenomegaly assay (Dietert et al. 1985). There was also a depressed cutaneous basophil hypersensitivity to injected PHA-P.

Considering that there are variations among the results of the studies on aflatoxin-impaired immunogenesis, there are generalized statements that have (Pier and McLoughlin 1985; CAST 1989) and can be made. The aflatoxins impair immunity without suppression of antibody synthesis, suppress non-specific humoral factors involved in immunogenesis, inhibit phagocytosis by macrophages, cause thymic aplasia, and suppress cell-mediated immunity.

Summary

Aflatoxins remain as a threat to the health of livestock as well as humans by their continuing intermittent occurrence in both feeds and foods. The finding that aflatoxin-contaminated feeds, and eventually purified aflatoxins, were carcinogenic in rats and trout initiated a multitude of studies in search of the role of these toxins in human liver disease, especially cancer. Although aflatoxins have caused acute liver disease in humans, epidemiologic evidence of the involvement of aflatoxins in PLC has not been clarified. Earlier studies did not consider that the hepatitis B virus (HBV) may have contributed to the PLC in the selected populations. Although later studies that did include measurement of the HBV antigen in serum provided conflicting evidence for the role of aflatoxin in PLC in these populations, the latest and most comprehensive study found no association between aflatoxin exposure and PLC mortality. The technological advances and findings of the chemical, immunologic, and metabolic activities of aflatoxins such as binding to DNA and protein to form adducts, development of monoclonal antibodies, and mutational specificity of the genotoxic compounds will, it is hoped, help to clarify the role of aflatoxin as a risk factor, among many others, in the development of primary liver cancer in humans.

Aflatoxicosis of animals is usually manifested by pathologic changes in the liver, but they have been found to be carcinogenic and teratogenic as

well as causing impaired protein formation, coagulation, weight gains, and immunity. The importance of the carcinogenic effect in livestock is diminished because they are not fed contaminated diets for a sufficient time prior to marketing for slaughter. Animals are variably susceptible to aflatoxins, depending on such factors as age, species, breed, sex, nutrition, and certain stresses.

Swine, cattle, and poultry are the domestic species of greatest economic concern in terms of aflatoxicosis. In all species, the evidence of disease is a general unthriftiness and reduction in weight gains, feed efficiency, immunity, and production. More conclusive evidence of aflatoxin involvement in disease includes acute to chronic liver disease with concomitant increases in specific liver enzymes in the serum. In cattle, milk production is affected, but of greater significance is that the aflatoxins in feeds can be rather efficiently converted to toxic metabolites in milk, with even small amounts being readily detectable. The poultry industry probably suffers greater economic loss than any of the livestock industries because of the greater susceptibility of their species to aflatoxins than other species.

Because other mycotoxins may occur in feeds in concert with aflatoxins, a consideration of their possible interactions in producing disease is important. The aflatoxins have been shown to interact synergistically with both T-2 toxin and ochratoxin in their effects on poultry.

Immunosuppression is a major economic effect of aflatoxins. Although antibody production is impaired when high concentrations of aflatoxins are consumed, the major effects are related to cellular immune phenomena or the reduction of nonspecific humoral factors such as complement and interferon. They cause thymic aplasia and inhibit phagocytosis by macrophages, delayed cutaneous hypersensitivity, graft vs host response, and interfere with lymphoblastogenesis and leukocyte migration. The effects of aflatoxins on infectious diseases are dependent on the disease agent, toxin dose and constitution, animal species, and perhaps sensitivity of the test. The embryonic exposure of animals to aflatoxin occurs and can affect the immune status of a neonatal animal.

References

- Adamson RH, Correa P, Dalgaard DW (1973) Brief communication: Occurrence of a primary liver carcinoma in a Rhesus monkey fed aflatoxin B₁. *J Natl Cancer Inst* 50:549-553
- Adinarayanaiah GL, Nanjiah RD, Keshavamurthy BS (1973) Effect of aflatoxin on antibody production to *Salmonella pullorum* antigen in chicks. *Indian Vet J* 2:297.
- Allcroft R (1965) Aspects of aflatoxicosis in farm animals. In: Wogan GN (ed) *Mycotoxins in Foodstuffs: Proceedings of MIT Symposium*, March 18-19, 1964. Cambridge, MA, pp 153-162.

- Alpert ME, Hutt MSR, Davidson CS (1969) Primary hepatoma in Uganda, a prospective clinical and epidemiological study in 46 patients. *Am J Med* 46:794–802.
- Applebaum RS, Brackett RE, Wiseman DW, Marth EH (1982) Responses of dairy cows to dietary aflatoxin: Feed intake and yield, toxin content, and quality of milk of cows treated with pure and impure aflatoxin. *J Dairy Sci* 65:1503–1508.
- Ambrecht BH, Wiseman HG, Shalkop WT (1972) Swine aflatoxicosis II. The chronic response in brood sows fed sublethal amounts of aflatoxin and the reaction in their piglets. *Environ Phys Biochem* 2:77–85.
- Asao T, Buche G, Abdel Kader MM, Chang SB, Wick EL, Wogan GN (1965) The structures of Aflatoxin B₁ and G₁. In: Wogan GH (ed) *Mycotoxins in Foodstuffs: Proceedings of MIT Symposium*, March 18–19, 1964. MIT Press, Cambridge, MA, pp 265–273.
- Astrup H, Bradley KA, Shamsuddin AKM, Wakhisi J, Wasunna A (1983) Detection of putative adduct with fluorescence characteristics identical to 2,3-dihydro-2(7-N-guanyl)-3-hydroxy-aflatoxin B₁ in human urine in Murang's District, Kenya. *Carcinogenesis* 9:1193–1195.
- Astrup H, Seremet T, Wakhisi J, Wasunna A (1987) Aflatoxin exposure measured by urinary excretion of aflatoxin B₁ guanine adduct and hepatitis B virus infection in areas with different liver cancer incidence in Kenya. *Cancer Res* 47:3420.
- Baertschi SW, Raney KD, Stone MP, Harris TM (1988) Preparation of 8,9 epoxide of the mycotoxin aflatoxin, B₁, the ultimate carcinogenic species. *J Am Chem Soc* 110:7929–7931.
- Bassendine MF (1987) Aetiological factors in hepatocellular cancer. *Bailliere's Clin Gastroenterol* 1(1):1–16.
- Blount WP (1960) A new turkey disease problem in England. *Brit Oil and Cake Mills, Ltd. Quar Poult Bull* 27:1–3.
- Blount WP (1961) Turkey "X" disease. *Turkeys* 9:2:52, 55, 58, 61, 77.
- Bodine AB, Mertens DR (1983) Toxicology, metabolism, and physiological effects of aflatoxins in the bovine. In: Diener UL, Asquith RL, Dickens JW (eds) *Aflatoxin and Aspergillus flavus in corn*. Alabama Agric Exp Sta, Auburn Univ, AL, pp 46–50.
- Boonchuvit B, Hamilton PB, Burneister HR (1975) Interaction of T₂ toxin w/salmonella infections in chickens. *Poult Sci* 54:1693–1696.
- Bressac B, Kew M, Wands J, Ozturk M (April 1991) Selective G to T mutations of p53 gene in hepatocellular carcinoma from southern Africa. *Nature* 350:429–430.
- Brown RW, Pier AC, Richard JL, Krogstad RE (1981) Effects of dietary aflatoxin on existing bacterial intramammary infections in dairy cattle. *Am J Vet Res* 42:927–933.
- Burg WR, Shotwell OL, Saltzman BE (1981) Measurements of airborne aflatoxins during the handling of contaminated corn. *Am Ind Hyg Assoc J* 42:1–11.
- Burg WR, Shotwell OL, Saltzman BE (1982) Measurements of airborne aflatoxins during the handling of 1979 contaminated corn. *Am Ind Hyg Assoc J* 43:580–586.
- Burg WR, Shotwell OL (1984) Aflatoxin levels in airborne dust generated from contaminated corn during harvest and at an elevator in 1980. *J Assoc Offic Anal Chem* 67:2:309–312.

- Busby WF Jr, Wogan GN (1981) Aflatoxins. In: Shank RC (ed) *Mycotoxins and N-Nitroso Compounds*. CRC Press, Boca Raton, FL, pp 3–27.
- Butler WH, (1965) Liver Injury and Aflatoxin. In: Wogan GH (ed) *Mycotoxins in Foodstuffs, Proceedings of MIT Symposium*, March 18–19, 1964, MIT Press, Cambridge, MA, pp 175–186.
- Byong IY, Rothenbacher H (1982) Immunopathologic effects of aflatoxin on lymphoid tissues and on the pathogenesis of Newcastle disease. *J Am Vet Med Assoc* 181:286 (abstract)
- Campbell ML Jr, May JD, Huff WE, Doerr JA (1983) Evaluation of immunity of young broiler chickens during simultaneous aflatoxicosis and orhratoxicosis. *Poult Sci* 62:2138–2144.
- Campbell TC, Caedo JP, Bullatto JJ, Salamet L, Engel RW (1970) Aflatoxin M₁ in human urine. *Nature (London)* 227:403–404.
- Campbell TC, Stoloff L (1974) Implication of mycotoxins for human health. *J Agr Food Chem* 22(6):1006–1015.
- Campbell TC, Chen J, Liu C, Li J, Parpia B (1990) Nonassociation of aflatoxin with primary liver cancer in a cross-sectional ecological survey in the People's Republic of China. *Cancer Res* 50:6882–6893.
- Carnaghan RBA (1965) Hepatic tumors in ducks fed a low level of toxic groundnut meal. *Nature* 203:308.
- CAST (1989) *Mycotoxin, Economic and Health Risks*. Task Force Rept 116.
- Caster, WO, Burton, TA, Irvin TR, Tanner MA (1986) Dietary aflatoxins, intelligence and school performance in Southern Georgia. *Int J Vit Nutr Res* 56:291–295.
- Cheng-ya W, Xin-sheng Y, Jin-Xin H, Tao-zhen L, Yao-qin Y (1987) HLA related genetic control of natural killer activity and aflatoxin B₁ suppression of lymphocytic blastogenesis. *Chin Med J Engl* 100:29–33.
- Cole RJ (1986) Etiology of turkey "X" disease in retrospect: A case for the involvement of cyclopiazonic acid. *Mycotoxin Res* 2:3–7.
- Cook WO, Richard JL, Osweiler GD, Trappel DW (1986) Clinical and pathologic changes in acute bovine aflatoxicosis: Rumen motility and tissue and fluid concentrations of aflatoxin B₁ and M₁. *Am J Vet Res* 47:1817–1825.
- Cook WO, Van Alstine WG, Osweiler GD (1989) Aflatoxins in Iowa swine: Eight cases (1983–1985) *JAVMA* 194:554–558.
- Cysewski SJ, Pier AC, Engstrom GW, Richard JL, Dougherty RW, Thurston JR (1968) Chemical pathologic features of acute aflatoxicosis of swine. *Am J Vet Res* 29:1577–1590.
- Cysewski SJ, Wood RL, Pick AC, Baetz AL (1978) Effects of aflatoxin on the development of acquired immunity to swine erysipelas. *Am J Vet Res* 39:445–448.
- Dalvi RR (1986) An overview of aflatoxicosis of poultry: Its characteristics, prevention and reduction. *Vet Res Comm* 10:429–443.
- Deger GE (1976) Aflatoxin—Human colon carcinogenesis? *Ann Intern Med* 85:204.
- Dietert RR, Qureshi MA, Nanna UC, Bloom SE (1985) Embryonic exposure to aflatoxin B₁: Mutagenicity and influence on development and immunity. *Environ Mut* 7:715–725.

- Doerr JA, Huff WE, Hamilton PB (1987) Increased sensitivity to staphylococcal beta hemolysins of erythrocytes from chickens during aflatoxicosis. *Poult Sci* 66:1929–1933.
- Duthie IF, Lancaster MC, Taylor J, Thomas DC, Shacklady CA, Attfield PH, Fuller-Lewis E (1966) Toxic groundnut meal in feeds for pig. I. A trial made at 2 laboratories with pigs from about 40–200 pounds live weight fed to a restricted scale. *Vet Rec* 79:621–625.
- Dvorackova I, Stora C, Ayraud N (1981) Evidence for aflatoxin B₁ in two cases of lung cancer in man. *Cancer Res Clin Oncol* 100:221–224.
- Dvorak R, Jagos J, Bouda J, Piskac A, Zapletal O (1977) Changes in the clinico-biochemical indices in the rumen liquor and urine in cases of experimental aflatoxicosis in dairy cows. *Vet Med (Prague)* 22:161–169.
- Edds GT, Nair KPC, Simpson CF (1973) Effect of aflatoxin B₁ on resistance in poultry against cecal coccidiosis and Marek's disease. *Am J Vet Res* 34:819–826.
- Essigmann JM, Croy RG, Nadzan AM, Busby WF Jr, Reinhold VN, Biechi G, Wogan GN (1977) Structural identification of the major DNA adduct formed by aflatoxin B₁ *in vitro*. *Proc Nat Acad Sci USA* 74:1870–1874.
- Fehr PM, Delage J (1970) Effect de l'aflatoxine sur les fermentations du rumen. *Can Nutr Diet* 5:59–61.
- Galikeev KHL, Raipov OR, Manyasheva RA (1968) Effect of aflatoxin on dynamics of antibody formation. *Byul Eksper Biol Medoc (USSR)* 65:88 (translation).
- Garlich JD, Tung HT, Hamilton PB (1973) The effects of short term feeding of aflatoxin on egg production and some plasma constituents of the laying hen. *Poult Sci* 52:2206–2211.
- Garner C, Ryder R, Montesano R (1985) Monitoring of aflatoxins in human body fluids and application to field studies. *Cancer Res* 45:922–928.
- Giambrone JJ, Ewert DL, Wyatt RD, Edison CS (1978) Effect of aflatoxin on the humoral and cell mediated immune systems of the chicken. *Am J Vet Res* 39:305–308.
- Goldblatt LA (ed) (1969) *Aflatoxin Scientific Background Control and Implications*. Academic Press, New York.
- Gopalakrishnan S, Harris TM, Stone MP (1990) Intercalation of aflatoxin B in two oligodeoxynucleotide adducts: Comparative ¹H NMR analysis of d(ATC^{AFB}GAT), d(ATCGAT) and d(AT^{AFB}GCAT)₂. *Biochemistry* 29:10438–10448.
- Groopman JD, Cain LG, Kensler TW (1988) Aflatoxin exposure in human populations: Measurements and relationship to cancer. *CRC Crit Rev Toxicol* 19:113–145.
- Groopman JD, Donohue K (1988) Aflatoxin, a human carcinogen: Determination in foods and biological samples by monoclonal antibody affinity chromatography. *J Assoc Anal Chem* 71:5:861–867.
- Groopman JD, Sabbioni G (1991) Detection of aflatoxin and its metabolites in human biological fluids. In: Bray GA, Ryan DH (eds) *Mycotoxins, Cancer, and Health*. Louisiana State Univ. Press, Baton Rouge, LA, pp 18–31.
- Guthrie LD (1979) Effects of aflatoxin in corn on production and reproduction in dairy cattle. *J Dairy Sci* 62:134.

- Halver JE (1969) Aflatoxicans and trout hepatoma. In: Goldblatt LA (ed) *Aflatoxin*. Academic Press, New York, pp 265–306.
- Hamilton PB (1971) A natural and extremely severe occurrence of aflatoxicosis in laying hens. *Poult Sci* 50:1880–1882.
- Hamilton PB, Garlich JD (1971) Aflatoxin as a possible cause of fatty liver syndrome in laying hens. *Poult Sci* 50:800–804.
- Hamilton PB, Harris JR (1971) Interaction of aflatoxins with *Candida albican* infections and other stresses in chicken. *Poult Sci* 50:906–912.
- Harris AL (April 1991) Telling changes of base. *Nature* 350:377–378.
- Hartley RD, Nesbitt BF, O'Kelley J (1963) Toxic metabolites of *Aspergillus flavus*. *Nature* 198:1056–1058.
- Hayes RB, Van Nieuwenhuize JP, Raatgever JW, ten Kate FJW (1984) Aflatoxin exposures in the industrial setting: An epidemiological study of mortality. *Food Chem Toxicol* 22:1:39–43.
- Hendrickse RG, Coulter JBS, Lamplugh SM, MacFarlane SBJ, Williams TE, Omer MIA, Sulimen GI, El-Zorgani GA (1983) Aflatoxins and kwashiorkor: Epidemiology and clinical studies in Sudanese children and findings in autopsy liver samples from Nigeria and South Africa. *Bull Soc Pathol Exp* 76:559–566.
- Hendrickse RG (1984) Kwashiorkor: 50 years of myth and mystery. Do aflatoxins provide a clue? *Acta Leidensia* 53:11–30.
- Hendrickse RG (1985) The influence of aflatoxins on child health in the tropics with particular reference to kwashiorkor. *Trans Royal Soc Trop Med Hyg* 78:427–435.
- Hintz HF, Heitman HJ, Booth AN, Gagne WE (1967) Effects of aflatoxin on reproduction in swine. *Proc Soc Exp Biol Med* 126:146–148.
- Hsu IC, Metcalf RA, Sun T, Welsh JA, Wang NJ, Harris CC (April 1991) Mutational hotspot in the p53 gene in human hepatocellular carcinomas. *Nature* 350:427–428.
- Huff WE, Doerr JA (1981) Synergism between aflatoxin and ochratoxin A in broiler chickens. *Poult Sci* 60:550–555.
- Huff WE, Doerr JA, Wabeck CJ, Chaloupka GW, May JD, Merkley JW (1983) Individual and combined effects of aflatoxin and ochratoxin A on bruising in broiler chickens. *Poult Sci* 62:1764–1771.
- Huff WE, Kubena LF, Harvey RB, Hagler WM, Jr, Swanson SP, Phillips TD, Creger CR (1986) Individual and combined effects of aflatoxin and deoxynivalenol (DON, vomitoxin) in broiler chickens. *Poult Sci* 65:1291–1298.
- Huff WE, Harvey RB, Kubena LF, Rottinghaus GE (1988) Toxic synergism between aflatoxin and T-2 toxic in broiler chickens. *Poult Sci* 67:1418–1423.
- International Agency for Research in Cancer (IARC) (1976) *Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man: Some Naturally Occurring Substances*, Vol 10. Lyon, France, pp 51–72.
- IARC (1987) *Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans: Aflatoxins*, Suppl 7. Lyon, France, pp 83–87.
- Joens LA, Pier AC, Cutlip R (1981) Effects of aflatoxin consumption on the clinical course of swine dysentery. *Am J Vet Res* 42:1170–1172.
- Kratzer FH, Bandy D, Wiley M, Booth AN (1968) Aflatoxin effects in poultry. *Proc Soc Exp Biol Med* 131:1281–1284.

- Kraybill HF, Shimkin MB (1964) Carcinogenesis related to foods contaminated by processing and fungal metabolites. *Adv Cancer Res* 8:191–248.
- Krishnamachari, KAVR, Bhat, RV, Nagarajan, V, Tilac, TBG (1975a) Investigations into an outbreak of hepatitis in Western India. *Ind J Med Res* 63:1036–1048.
- Krishnamachari, KAVR, Bhat, RV, Nagarajan, V, Tilac, TBG (1975b) Hepatitis due to aflatoxicosis. *Lancet* 1:1061–1063.
- Lancaster MC, Jenkins FP, Philp JM (1961) Toxicity associated with certain samples of groundnuts. *Nature (London)* 192:1095–1096.
- Lancaster MC (1968) Comparative aspects of aflatoxin-induced hepatic tumors. *Cancer Res* 28:2288–2292.
- Larsen AB, Cysewski SL, Miller JM (1975) Effects of aflatoxin on susceptibility of hamsters to mycobacterium paratuberculosis. *Am J Vet Res* 36:1545–1547.
- Masri MS, Garcia VC, Page JR (1969) The aflatoxin M₁ content of milk from cows fed known amounts of aflatoxin. *Vet Rec* 84:146–147.
- McLoughlin ME, Pier AC, Thurston JR (1984) Use of bacteria and yeasts to identify T lymphocytes in peripheral blood and lymphoid tissues of healthy guinea pigs fed aflatoxin. *Am J Vet Res* 45:1795–1799.
- Mertens DR (1979) Biological effects of mycotoxins upon rumen function and lactating dairy cows. In: *Proceedings of Symposium on Interaction of Mycotoxins in Animal Production*, Michigan State Univ, Lansing, July 13, 1978. *Natl Acad Sci, Washington, DC*, pp 118–136.
- Michael GY, Thaxton P, Hamilton PB (1973) Impairment of the reticuloendothelial system of chickens during aflatoxicosis. *Poult Sci* 52:1206–1207.
- Miller DM, Stuart BP, Crowell WA (1981) Experimental aflatoxicosis in swine. Morphological and clinical pathological results. *Can J Corp Med* 45:343–351.
- Miller DM, Crowell WA, Stuart BP (1982) Acute aflatoxicosis in swine: Clinical pathology, histopathology, and electron microscopy. *Am J Vet Res* 43:273–277.
- Ngindu A, Kenya PR, Ocheng DM, Omonde TN, Ngare W, Gatei D, Johnson BK, Ngira, JA, Nandwa, H, Jansen, AJ, Kaviti, JN, Siongok, TA (1982) Outbreak of acute hepatitis caused by aflatoxin poisoning in Kenya. *Lancet* 1:1346–1348.
- Oettle AG (1965) The etiology of primary carcinoma of the liver in Africa: A critical appraisal of previous ideas with an outline of the mycotoxin hypothesis. *S. A. Tydskrif Vir Genees-kunde* 9 October:817–825.
- Panangala VS, Giambrone JJ, Diener UL, Davis ND, Hoerr FJ, Mitra A, Schultz RD, Wilt GR (1986) Effects of aflatoxin on the growth performance and immune responses of weanling swine. *Am J Vet Res* 47:2062–2067.
- Park DL, Pohland AE (1986) A rationale for the control of aflatoxin in animal feeds. In: Steyn PS, Vleggaar R (eds) *Mycotoxins and Phycotoxins*. Presented at the 6th International IUPAC Symposium on Mycotoxins and Phycotoxins, Pretoria, Republic of South Africa, July 22–25, 1985. Elsevier Scientific Pub, Amsterdam, The Netherlands, pp 473–482.
- Patterson DSP, Anderson PH (1982) Recent aflatoxin feeding experiments in cattle. *Vet Rec* 110:60.
- Peers FG, Linsell CA (1973) Dietary aflatoxins and liver cancer—a population study in Kenya. *Brit J Cancer* 27:473–484.
- Peers FG, Gilman CA, Linsell CA (1976) Dietary aflatoxins and human liver cancer—a study in Swaziland. *Int J Cancer* 17:167–176.
- Peers F, Bosch X, Kaldor J, Linsell A, Pluijmen M (1987) Aflatoxin exposure, hepatitis B virus infection and liver cancer in Swaziland. *Int J Cancer* 39:545–553.
- Pier AC, Heddleston KL (1970) The effect of aflatoxin on immunity on turkeys. I. Impairment of actively acquired resistance to bacterial challenge. *Avian Dis* 14:797–809.
- Pier AC, Heddleston KL, Boney WA, Lukert PD (1971) The effect of aflatoxin on immunity. *Proc XIX Cong Mundial de Med Vet Zootech* 1:216–219.
- Pier AC, Cysewski SJ, Richard JL, Baetz AL, Mitchell L (1976) Experimental mycotoxicosis in calves with aflatoxin, ochratoxin, rubratoxin, and T-2 toxin. In: *Proceedings 80th Annual Meeting of US Animal Health Association*, pp 130–148.
- Pier AC (1981a) Mycotoxins and animal health. *Adv Vet Sci Comp Med* 25:185–243.
- Pier AC (1981b) Mycotoxins and animal health. *Adv Vet Sci Comp Med* 25:185–243.
- Pier AC, McLoughlin ME, Richard JL, Baetz A, Dahlgren RR (1985) *In utero* transfer of aflatoxin and selected effects in neonatal pigs. In: Lacey J (ed) *Trichothecenes and Other Mycotoxins*. Wiley, New York, pp 495–506.
- Pier AC, McLoughlin ME (1985) Mycotoxic suppression of immunity. In: Lacey J (ed) *Trichothecenes and Other Mycotoxins*. Wiley, New York, pp 507–519.
- Pier AC (1986) Immunomodulation in aflatoxicosis. In: Richard JL, Thurston JR (eds) *Diagnosis of Mycotoxicoses*. Martinus Nijhoff Pub, Dordrecht; pp 143–148.
- Price RL, Paulson JH, Lough OG, Gingg C, Kurtz AG (1984) Aflatoxin conversion by dairy cattle consuming naturally contaminated whole cottonseed. *Bull Dept Nutr Food Sci Univ Arizona, Tucson, AZ*.
- Ratanesthankul C (1983) Effect of aflatoxin on immune response to Australian infectious bronchitis virus in chickens. Ph.D. diss, Univ. Sydney, Australia.
- Reddy RV, Taylor MJ, Sharma RP (1987) Studies of immune function of CD-1 mice exposed to aflatoxin B₁. *Toxicol* 43:123–132.
- Reye RDK, Morgan G, Baral J (1963) Encephalopathy and fatty degeneration of the viscera: A disease entity in childhood. *Lancet* 2:749–752.
- Richard JL, Pier AC, Cysewski SJ, Grehan CK (1973) Effect of aflatoxin and aspergillosis on turkey poults. *Avian Dis* 17:111–121.
- Richard JL, Thurston JR (1975) Effect of aflatoxin on phagocytosis of *Aspergillus fumigatus* spores by rabbit alveolar macrophages. *Appl Microbiol* 30:44–77.
- Richard JL, Thurston JR, Pier AC (1978) Effects of mycotoxins on immunity. In: Rosenberg P (ed) *Toxins: Animal, Plant, and Microbial*. Pergamon Press, New York, pp 801–817.
- Richard JL, Pier AC, Stubblefield RD, Shotwell OL, Lyon RL, Cutlip RC (1983) Effect of feeding corn naturally contaminated with aflatoxin on feed efficiency, on physiologic, immunologic, and pathologic changes, and on tissue residue in steers. *Am J Vet Res* 44:1294–1299.
- Richard JL, Stubblefield RD, Lyon RL, Peden WM, Thurston JR, Rimler RB (1986) Distribution and clearance of aflatoxins B₁ and M₁ in turkeys fed diets containing

- 50 or 150 ppb aflatoxins from naturally contaminated corn. *Avian Dis* 30:788-793.
- Richard JL (1991) Mycotoxins as immunomodulators in animal systems. In: Bray GA, Ryan DH (eds) *Mycotoxins, Cancer, and Health*. Pennington Nutrition Series, Louisiana State Univ Press, Baton Rouge, LA, pp 197-220.
- Rogan WJ, Yang GC, Kimbrough RD (1985) Aflatoxin and Reye's syndrome: A study of livers from diseases cases. *Arch Environ Hlth* 40:2:91-95.
- Ryan NJ, Hogan GR, Hayes AW, Unger PD, Siraj MY (1979) Aflatoxin B₁: Its role in the etiology of Reye's syndrome. *Pediatrics* 64(1):71-75.
- Sabbioni G, Skipper PL, Buchi G, Tannenbaum SR (1987) Isolation and characterization of the major serum albumin adduct formed by aflatoxin B₁ *in vivo* in rats. *Carcinogenesis* 8(6):819-824.
- Sargeant K, Shendan A, O'Kelly J, Carnahan RBA (1961) Toxicity associated with certain samples of groundnuts. *Nature* 192:1096-1097.
- Shalkop WT, Armbrrecht BH (1974) Carcinogenic response of brood sows fed aflatoxin for 28 to 30 months. *Am J Vet Res* 35:623-627.
- Shank RC, Gordon JE, Wogan GN (1972) Dietary aflatoxins and human liver cancer. III. Field survey of rural Thai families for ingested aflatoxins. IV. Incidence of primary liver cancer in two municipal populations of Thailand. *Food Cosmet Toxicol* 10:71-84, 171-179.
- Shank RC (1977) Epidemiology of aflatoxin carcinogenesis. *Adv Mod Toxicol* 3:291-318.
- Sherlock S, Fox RA, Scheuer PJ (1970) Chronic liver disease and primary liver-cell cancer with hepatitis-associated (Australia) antigen in serum. *Lancet* 1:1243-1247.
- Shimada T, Guengrich FP (1989) Evidence for cytochrome P-450-N-F the nifedipine oxidase being the principal enzyme involved in the bioactivation of aflatoxins in human liver. *Proc Natl Acad Sci* 86:462-465.
- Silas JC, Harrison MA, Carpenter JA, Roth IL (1987) Airborne aflatoxin in corn processing facilities in Georgia. *Am Ind Hyg Assoc J* 48(3):198-201.
- Sisk DB, Carlton WW (1972) Effect of dietary protein concentration on response. *Am J Vet Res* 33:107-114.
- Smith JW, Prince WR, Hamilton PB (1969) Relationship of aflatoxins to *Salmonella gallinarum* infections of chickens. *Appl Microbiol* 18(5):946-947.
- Southern LL, Clawson AJ (1979) Effects of aflatoxins on finishing swine. *J An Sci* 49:1006-1011.
- Stoloff L (1977) Aflatoxins—An overview. In: *Mycotoxins in Human and Animal Health*. Rodrick, Hesseltine and Mehlman, Pathotox, pp 7-8.
- Stoloff L (1983) Aflatoxin as a cause of primary liver-cell cancer in the United States: A probability study. *Nutr Cancer* 5:3-4,165-185.
- Sun T-T, Chu Y-Y (1984) Carcinogenesis and prevention strategy of liver cancer in areas of prevalence. *J Cell Physiol* 3(Suppl.):39-44.
- Sun TT, Wu SM, Wu YY, Chu YR (1986) Measurement of individual aflatoxin exposure among people having different risk to primary hepatocellular carcinoma. In: Hayashi Y et al. (eds), *Diet, Nutrition and Cancer*. Japan Sci Soc Press, Tokyo/VNU Sci Press, Utrecht, pp 225-235.

- Thaxton JP, Tung HT, Hamilton PB (1974) Immunosuppression in chickens by aflatoxin. *Poult Sci* 53:721.
- Thurston JR, Richard JL, Cysewski SJ, Pier AC, Graham CK (1972) Effect of aflatoxin on complement activity in guinea pigs. *Proc Soc Exp Biol Med* 138:300-303.
- Thurston JR, Deyoe BL, Baetz AL, Richard JL, Booth GD (1974) Effect of aflatoxin on serum proteins, complement activity, and the antibody response to *Brucella abortus* in guinea pigs. *Am J Vet Res* 35:1097.
- Thurston JR, Cook W, Driftmier K, Richard JL (1986). Decreased complement and bacteriostatic activities in the sera of cattle given single or multiple doses of aflatoxin. *Am J Vet Res* 47:836-849.
- Truckess MW, Richard JL, Stoloff L, McDonald JS, Brumby WC (1983) Absorptions and distribution patterns of aflatoxins B₁ and M₁ in blood and milk of cows given aflatoxin B₁. *Am J Vet Res* 44:1753-1756.
- Van Dijk HJ, O'Dell GD, Bodine AB (1984) Effects of aflatoxin M₁ intake, its physiologic levels on newborn dairy calves. *Am J Vet Res* 45:1994-1997.
- Van Nieuwenhuize JP, Herber RFM, de Bruin A, Meyer PB, Duba WC (1973) Aflatoxinen: Epidemiologisch onderzoek naar carcinogeniteit bij langdurige "low level" expositie van een fabriekspopulatie. *T. Soc Geneesk* 51:754.
- Van Rensburg SJ, Van Der Watt JJ, Purchase IFH, Pereira Coutinho L, Markham L (1974) Primary liver cancer rate and aflatoxin intake in a high cancer area. *S Afr Med J* 48:2508a-2508d.
- Van Rensburg SJ, Cook-Mozaffari P, Van Schalkwyk DJ, Van Der Watt JJ, Vincent TJ, Purchase IF (1985) Hepatocellular carcinoma and dietary aflatoxin in Mozambique and Transkei. *Brit J Cancer* 51:713-726.
- Wagstaff DJ (1985) The use of epidemiology, scientific data, and regulatory authority to determine risk factors in cancers of some organs of the digestive tract. 3. Liver cancer. *Reg Toxicol Pharmacol* 5:384-404.
- Wagstaff DJ (1990) Epidemiologic studies of the association of aflatoxin exposure and human liver cancer. In: *Proceedings of Symposium on Perspective on Aflatoxin in Field Crops and Animal Food Products in the United States*, Washington, DC, Jan 23-24, 1990. ARS-83; pp 141-147.
- Wilson, BJ (1978) Hazards of mycotoxins to public health. *J Food Protect* 41:5:375-384.
- Wogan GN, Newberne PM (1967) Dose-response characteristics of aflatoxin B₁ carcinogenesis in the rat. *Cancer Res* 27:2370.
- Wolzak A, Pearson AM, Coleman TH, Pestka JJ, Gray JI (1985) Aflatoxin deposition and clearance in the eggs of laying hens. *Food Chem Toxicol* 23:1057-1061.
- Wyatt RD, Ruff MD, Page RK (1975) Interaction of aflatoxin with *Elmera tenella* infection and monensin in young broiler chickens. *Avian Dis* 19(4):730-740.
- Yang WC (1983) Effects of aflatoxin B₁ on the development of porcine cellular and humoral immune responses. *Dissertation Abs Int* 44/06-B:1778.
- Yeh FS, Yu MC, Mo CC, Luo S, Tong MJ, Henderson BE (1989) Hepatitis B virus, aflatoxins, and hepatocellular carcinoma in southern Guangxi, China. *Cancer Res* 49:2506-2509.

Zhu JQ, Zhang LS, Hu X, Xiao Y, Chen JS, Xu YC, Fremy J, Chu FS (1987)
Correlation of dietary aflatoxin B1 levels with excretion of aflatoxin M1 in human
urine. *Cancer Res* 47:1848–1852.

Manuscript received December 18, 1991; accepted January 20, 1992.